



Guidelines for Adequate Water Intake : A Public Health Rationale



Proceedings from EFBW Symposium
IUNS 20th International Congress of Nutrition
Granada, Spain, September 18, 2013

Over the past years, the European Food Safety Authority (EFSA) has issued two important scientific opinions related to Water. In March 2010, EFSA released its Scientific Opinion on Dietary Reference Values for Water, and in April 2011 EFSA released its Scientific Opinion on water related health claims.

These Scientific opinions were based on an extensive review of scientific evidence. They are extremely important because they clarify the reference values for water among the population in Europe according to age or physiological status (Table1) and confirm the importance of water for the maintenance of basic physiological functions such as thermoregulation, physical and cognitive function. To put it simply, they highlight the importance of drinking enough water (2 litres per day from food and drink) to allow our body to perform optimally.

Table 1. Dietary Reference Values for water EFSA (2010)			
		Total water adequate intake	
Infants	0-6 months	680 mL/d (through milk)	
	6-12 months	800-1000 mL/d	
Children	1-2 years	1100-1200 mL/d	
	2-3 years	1100-1200 mL/d	
	4-8 years	1600 mL/d	
	9-13 years	Boys	2100 mL/d
		Girls	1900 mL/d
> 14 years	Cf adults		
Adults	Men	2500 mL/d	
	Women	2000 mL/d	
Pregnant women		+ 300 mL/d vs adults	
Lactating women		+ 600-700 mL/d vs adults	
Elderly		Same as adults	

EFSA also clarified the contribution of the water from food versus the water from drinks. Despite it being highly variable as it depends on the individual's diet for that day, food contributes on average 20% of the total water intake (EFSA, 2010). This means that on average, 80% of our total water intake should come from the fluid we drink. This gives values ranging from 1.6 L/D of water for women to 2.0 L/D of water for men via drinks. It is important to note that those values have to be adjusted depending on climate and physical activity. Exercising can easily increase water losses through sweat - up to one litre per hour or more. This quantity of water lost by the body has to be compensated through water intake on top of EFSA's Dietary reference values.

Despite variations between Member States, data shows that on average, in most European countries, a significant part of the population is actually drinking less than the values recommended by EFSA. In the interest of those who are not drinking enough, it would therefore be beneficial to promote EFSA's dietary reference values more actively at national level.

The symposium *Adequate Water Intake – Public Health Rational* organised by the European Federation of Bottled Waters (EFBW) gathers International Scientific Experts in their field who will present the science supporting EFSA's scientific opinions.



Dr. Laurent Le Bellego
Chairman of the
Health Group of the EFBW

Establishing International Guidelines for Adequate Water Intake

Lawrence E. Armstrong, Ph.D., FACSM, USA

Water is the matrix which supports digestion, metabolism, nutrient transport, cardiovascular function, and temperature regulation. However, because human activities are very diverse and because body water regulation is complex and dynamic, no consensus exists regarding the water requirements of adults, children, or special populations. This presentation will review the water Adequate Intake (AI) recommendations of nutritional and scientific organizations in Europe, China, Australia and the United States. These recommendations (a) include the total daily fluid intake of drinking water, beverages of all kinds, and food moisture; (b) distinguish youth, older age groups, and special populations such as pregnant women; and (c) apply to conditions of moderate environmental temperature and moderate physical activity levels. This presentation will review the factors that were considered, when each organization developed recommended AI values (i.e., cultural dietary customs, learned drinking habits, dental health, human performance, aging, and relationship of low daily water intake to chronic health outcomes). Further, the following factors will be presented as important considerations for interpreting AI values: water volume per unit of energy consumed, maximal safe water intake, and desirable urine osmolality.

References

- European Food Safety Authority (2011) Scientific opinion on dietary reference values for water. EFSA J 8, 1459–1506.
- Institute of Medicine (2004) Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate. Washington, DC: National Academies Press.
- Armstrong LE (2007) Assessing hydration status: the elusive gold standard. J Am Coll Nutr 26, 575S–584S.
- Ma G, Zhang Q, Liu A, Zuo J, Zhang W, Zou S, Li X, Lu L, Pan H, Hu X. Fluid intake of adults in four Chinese cities. Nutrition Reviews 70 (Suppl. 2): S105–S110, 2013.
- Department of Health & Ageing, National Health & Medical Research Council, Australia. Nutrient reference values for Australia and New Zealand. 2006.

European health claims: water intake is essential for the maintenance of normal physical function and thermoregulation - a scientific review.

Stavros A. Kavouras, PhD, FACSM, FECSS, USA

Water is the most essential constituent of the human body undergoing continuous recycling. Body water functions as a solvent, a means for heat balance, and a regulator of cell volume and overall functioning. Its balance in humans has been regulated to a large extent by the sensation of thirst; a key survival instinct in body fluid homeostasis^{1,2}.

For over a century, athletes have noticed that exercise-induced dehydration has been linked to decreased exercise performance. Francisco Lázaro, a Portuguese marathon runner was one of the first to notice that excessive sweating was making him slower. Therefore, he decided to stop dehydration by blocking sweating via application of wax over his skin during the 1912 Stockholm Olympics marathon. Unfortunately, he collapsed at the 30 km mark; and he died next morning from complications of hyperthermia. Since mid 40s, scientists have also been interested in the effect of dehydration on exercise performance. For instance, in 1944, Pitts and his colleagues showed that dehydration, as a response to exercise in the heat, increased thermoregulatory strain and induced exhaustion³.

Several years later, Armstrong et al examined the effect of hypohydration on exercise performance⁴. They studied the performance

of 8 trained, male runners on 1,500 m, 5,000 m, and 10,000 m while normally hydrated or hypohydrated on six separate occasions. Their running performance was reduced by 3.1, 6.7, and 6.3% in the 1,500 m, 5,000 m, and 10,000 m, respectively as a response to hypohydration. This decreased in exercise performance has been shown to be related to a significant decrease of stroke volume and cardiac output during exercise in the heat as a response to hypohydration. In the early 90s, Montain and Coyle concluded that athletes' increase in heart rate and core temperature, as well as, decline in stroke volume was proportional to the degree of hypohydration during exercise⁵. Similarly, Walsh and his colleagues, from the University of Cape Town in South Africa, found that even a mild decrease on hypohydration (-1.8% of body mass) decreased exercise performance; probably due to increased perception of fatigue, as indicated by the greater ratings of perceived exertion⁶. Even though that during exercise in the heat, hyperthermia and dehydration coincide it has been shown that both, hyperthermia and hypohydration, independently decreased athletes' ability to maintain cardiac output during exercise. Also hyperthermia and dehydration have an additive effect on the decline of cardiac output similar to the sum of the decline for each factor alone⁷. Even though hypohydration has been mainly studied during exercise in the heat, it

seems that even in a temperate environment hypohydration can limit exercise performance⁸.

In some sports events (ie, uphill cycling), athletes often believe that a decrease in body mass, even via dehydration, can enhance exercise performance by increasing the power to mass ratio. In this respect, Ebert et al. examined the effect of hypohydration during a hill cycling (8% grade) time exhaustion test in a warm environment at intensity equal to 88% of maximal aerobic power (MAP)⁹. Prior to the performance, test subjects cycled for 2 hour at 53% of their MAP, while drinking different volume of fluid in order to achieve the desired pre-exercise body mass (-2.5 vs. +0.3% of body mass). The results showed that even though cyclists were able to reduce their power requirements for any given cycling speed by lowering body mass; hypohydration had a significant detrimental effect on their performance. Hypohydration has been also linked to declined skill performance in sports like basketball, indicating the time to completion basketball-specific movement drill was slower and fewer shot were made when hypohydration reached -2% of body mass¹⁰. Recently, progressive dehydration induced great thermoregulatory and cardiovascular strain, as well as, whole body carbohydrate oxidation and muscle glycogenolysis¹¹. Interestingly, Bardis et al cycling speed and power output were lower in cyclist undergoing

dehydration of -1% of their body weight, due to higher body temperature and lower sweating sensitivity¹².

Compelling evidence supports the idea that hypohydration can decrease endurance exercise performance at least in hot and temperate environments, probably due to an increased thermoregulatory strain and cardiovascular decline. More recent data indicate that even a small degree of hypohydration, less than -2% of body mass, might decrease performance; thus, putting into question the current guidelines of the American College of Sports Medicine specifying that the goal of drinking during exercise should be to avoid dehydration greater than -2% of body mass.

References

- 1. Kavouras SA. Assessing hydration status. *Curr Opin Clin Nutr Metab Care*. Sep 2002;5(5):519-524.
- 2. Kavouras SA, Anastasiou CA. Water physiology. Essentiality, metabolism, and health implications. *Nutrition Today*. 2010;45(65):S27-S32.
- 3. Pitts G, Jonson R, Consolazio F. Work in the heat as affected by intake of water salt and glucose. *American Journal of Physiology*. 1944;142:253-259.
- 4. Armstrong LE, Costill DL, Fink WJ. Influence of diuretic-induced dehydration on competitive running performance. *Med Sci Sports Exerc*. Jul 31 1985;17(4):456-461.
- 5. Montain SJ, Coyle EF. Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise. *J Appl Physiol*. Sep 30 1992;73(4):1340-1350.
- 6. Walsh RM, Noakes TD, Hawley JA, Dennis SC. Impaired high-intensity cycling performance time at low levels of dehydration. *Int J Sports Med*. Oct 1994;15(7):392-398.
- 7. González-Alonso J, Mora-Rodríguez R, Below PR, Coyle EF. Dehydration markedly impairs cardiovascular function in hyperthermic endurance athletes during exercise. *J Appl Physiol*. Apr 31 1997;82(4):1229-1236.
- 8. Chevront SN, Carter R, Castellani JW, Sawka MN. Hypohydration impairs endurance exercise performance in temperate but not cold air. *J Appl Physiol*. Nov 2005;99(5):1972-1976.
- 9. Ebert TR, Martin DT, Bullock N, et al. Influence of Hydration Status on Thermoregulation and Cycling Hill Climbing. *Med Sci Sports Exerc*. Mar 01 2007;39(2):323-329.
- 10. Baker LB, Dougherty KA, Chow M, Kenney WL. Progressive dehydration causes a progressive decline in basketball skill performance. *Med Sci Sports Exerc*. Jul 30 2007;39(7):1114-1123.
- 11. Logan-Sprenger HM, Heigenhauser GJF, Killian KJ, Spriet LL. Effects of dehydration during cycling on skeletal muscle metabolism in females. *Med Sci Sports Exerc*. Oct 2012;44(10):1949-1957.
- 12. Bardis C, Kavouras S, Kosti L, Markousi M, Sidossis L. Mild Hypohydration Decreases Cycling Performance in the Heat. *Med Sci Sports Exerc*. 2013.

Water Intake is Essential for the Maintenance of Normal Cognitive function - a scientific review

Harris R. Lieberman, Ph.D., USA

Failure to maintain adequate hydration leads to a variety of adverse consequences, both physiological and psychological. The physiological effects of dehydration are well documented and are discussed by the other speakers at this symposium. The effects of mild or moderate levels of dehydration on human cognitive function are not as well recognized (Grandjean and Grandjean, 2007; Lieberman, 2007a). When humans or other animals are severely dehydrated the behavioral effects of this state are readily apparent - confusion, lethargy, and degraded cognitive function are present. In fact, in elderly individuals who are confused and disoriented, one of the first medical actions is assessing their state of hydration. Acute or chronic dehydration in elderly individuals can masquerade as much more serious and chronic medical conditions such as Alzheimer's disease (Wotton et al, 2008).

The assessment of cognitive state is a complex undertaking since there are so many different kinds of behavioral functions and a wide variety of methods exist to study them (Gawron, 2000; Lieberman, 2007b). Cognitive function includes a vast array of mental capabilities and states. Functions such as vigilance, learning, memory, reasoning, speech perception and hundreds of other activities are aspects of cognitive performance. Cognitive functions also include moods, symptoms and

emotions such as alertness, sadness, anxiety, and anger. Any of these may be sensitive to the effects of mild or moderate dehydration. However, only a small number of studies have been conducted to examine the effects of dehydration on cognitive function and only a few cognitive measures utilized in these studies. As with research on many stressors, most studies of dehydration have measured cognitive performance and mood. These studies clearly demonstrate that relatively high levels of dehydration, over 3% loss of body weight, impair both performance and mood. However, the effects of more moderate levels of dehydration are more controversial and the level of dehydration where cognitive function is first impaired has not been definitely established (Lieberman, 2007a). Several studies conducted in the 1980's at an Indian Defense Institute found that dehydration induced by exposure to high heat and exercise impaired specific aspects of cognitive function in healthy young men including short-term memory, reasoning and coordination. These deficits were apparent at 3% or higher levels of dehydration (Gopinathan et al, 1988; Sharma et al, 1986). Another more recent study, conducted with young male and female volunteers, induced a mean level of dehydration of 2.6% by 28 hours of fluid deprivation (Szinnai et al, 2005). This study found that mood states such as tiredness and concentration were degraded at this level of

dehydration but that cognitive performance was not affected, although females appeared more sensitive than males.

Recently, my colleagues at the University of Connecticut and I conducted two studies, using nearly identical methods, designed to assess the effects of mild dehydration on cognitive function (Armstrong et al, 2012; Ganio et al, 2011). One study was conducted with 26 young men and other with 25 young women. Dehydration was induced by moderate exercise in comfortable conditions. Both of these studies used a crossover procedure such that each individual was tested in a dehydrated and euhydrated state so each could serve as his/her own control. We assessed cognitive function using a comprehensive computerized battery of tests of cognitive performance, mood and symptoms associated with dehydration. The men in the study reached a 1.59% level of dehydration the women were 1.36% dehydrated on average. We found, in both men and women, that these levels of dehydration induced fatigue and degraded a specific aspect of cognitive function, vigilance. Women reported headache, confusion and lack of energy when mildly dehydrated and appeared somewhat more sensitive to the adverse effects of dehydration than men. These studies demonstrate that mild dehydration, which can occur over the course of daily activities, degrades cognitive function. They indicate how important it is to maintain adequate hydration by regularly consuming water so that optimal cognitive performance and mood are maintained throughout the day and adverse symptoms such as headache and fatigue are avoided.

References

- Armstrong LE, Ganio MS, Casa DJ, et al. Mild dehydration affects mood in healthy young women. *J Nutr.* 2012;142:382–388.
- Ganio MS, Armstrong LE, Casa DJ, et al. Mild dehydration impairs cognitive performance and mood of men. *Br J Nutr.* 2011;106:1535–1543.
- Gawron VJ. *Human Performance Measures Handbook.* Hillsdale, NJ: Lawrence Erlbaum Associates; 2000
- Gopinathan PM, Pichan G, Sharma VM. Role of dehydration in heat stress-induced variations in mental performance. *Arch Env Health.* 1988;43:15–17.
- Grandjean AC, Grandjean NR. Dehydration and cognitive performance. *J Am Coll Nutr.* 2007;26:549S–554S.
- Lieberman HR. Cognitive methods for assessing mental energy. *Nutr Neurosci.* 2007a;10:229–242.
- Lieberman HR. Hydration and cognition: a critical review and recommendation for future research. *J Am Coll Nutr.* 2007b;26:555S–561S.
- Sharma VM, Sridharan K, Pichan G, Panwar MR. Influence of heat-stress induced dehydration on mental functions. *Ergonomics.* 1986;29:791–799.
- Szinnai G, Schachinger H, Arnaud MJ, Linder L, Keller U. Effect of water deprivation on cognitive-motor performance in healthy men and women. *Am J Physiol Regul Integr Comp Physiol.* 2005;289:R275–R280.
- Wotton K, Crannitch K, Munt R. Prevalence, risk factors and strategies to prevent dehydration in older adults. *Contemp Nurse.* 2008;31:44–56.

About EFBW

EFBW is the voice of the European bottled water industry. A not-for-profit trade association with a membership base that includes national associations, companies and scientific bodies, EFBW collectively represents more than 600 producers of natural mineral water* and spring water* across Europe.

The Federation works to ensure that naturally sourced waters continue to offer a high quality, pure and convenient way to hydrate, and represents a sustainable and responsible choice for Europeans. The sector accounts for more than 158,000 direct employees throughout the production chain.

* Natural mineral waters and spring waters must both be from a designated underground origin only and be safe to drink at source, where they must be bottled directly. They may not be disinfected nor chemically treated.

In addition, natural mineral waters are characterised by their original purity and their stable mineral content as indicated on the label. In some cases, natural mineral waters can also be characterised by some health effects.



EFBW - European Federation of Bottled Waters aisbl

Rue de l'Association 32

BE-1000 Brussels

Tel.: + 32 2 210 20 32 Fax: + 32 2 210 20 35

www.efbw.eu